

- arteriogram report and a scoring system. *Clin Radiol* 1977; 28: 361-8
- 35 Nussmeier NA. Adverse neurological events: risks of intracardiac versus extracardiac surgery. *J Cardiothorac Vasc Anesth* 1996; 10: 31-7
- 36 Aberg T and Kihlgren M. Cerebral protection during open-heart surgery. *Thorax* 1977; 32: 525-33
- 37 Townes BD, Bashein G, Hornbein TF, et al. Neurobehavioural outcomes in cardiac operations: a prospective controlled study. *J Thorac Cardiovasc Surg* 1989; 98: 774-82
- 38 Wolman RL, Kanchuger MS, Newman ME, Roach GW and Nussmeier NA. Adverse neurologic outcome following intracardiac versus extracardiac surgery [Abstract]. *Perfusion* 1994; 9: 406
- 39 Murkin JM. Neurological dysfunction after CAB or valvular surgery: is the medium the miscreant? *Anesth Analg* 1993; 76: 213-4
- 40 Shaw PJ, Bates D, Cartlidge NEF, et al. Early intellectual dysfunction following coronary bypass surgery. *Q J Med* 1986; 58: 59-68
- 41 Aberg T, Ronquist G, Tyden H, et al. Adverse effects on the brain in cardiac operations as assessed by biochemical, psychometric, and radiologic methods. *J Thorac Cardiovasc Surg* 1984; 87: 99-105

An earlier version of this paper was presented by Dr Gorman at the SPUMS Annual Scientific Meeting 1998 in Palau.

Dr Simon Mitchell, MB ChB, Dip DHM, who is the Vice-President of the Underwater and Hyperbaric Medical Society, was Director of the Slark Hyperbaric Unit at the Royal New Zealand Navy Hospital, Naval Base, Auckland, New Zealand. This work was part of his PhD research program. He is now Medical Director of the Wesley Centre for Hyperbaric Medicine, Sandford Jackson Building, Chasely Street, Auchenflower, Queensland 4066, Australia. Telephone + 61-7-3371-6033. Fax 61-7-3371-1566. E-mail smitchell@wesley.com.au .

Ora Pellett, MSc, is a research psychologist at the University of Auckland, New Zealand. She is completing a PhD in Psychology in which she is studying the neurocognitive effects of cardiac surgery and their modification by protective strategies.

Professor D F Gorman FAFOM, PhD, is Head, Occupational Medicine, Faculty of Medicine and Health Sciences, University of Auckland, New Zealand. He is a Past-President of SPUMS. His address is 52 Albert Road, Devonport, Auckland 9, New Zealand. Telephone + 64-9-373-7599. Fax + 64-9-308-2379. E-mail d.gorman@auckland.ac.nz .

© 1999 by The Society of Thoracic Surgeons and reprinted, with minor changes, by kind permission of Dr Thomas B Ferguson, Editor of THE ANNALS OF THORACIC SURGERY, from Ann Thorac Surg 1999; 67: 1117-1124 where the title was "Cerebral Protection by Lidocaine During Cardiac Operations".

SPUMS ANNUAL SCIENTIFIC MEETING 1999

THE EFFECT OF BUBBLES ON THE LIVING BODY

Alf Brubakk

Key Words

Bubbles, decompression illness, physiology.

Introduction

There is general agreement that the basic problem in decompression is gas coming out of solution and forming a gas phase. However, it is also well known that a considerable number of bubbles can be formed without any acute signs or symptoms. Such bubbles have been called "silent" bubbles¹ and have, in particular, been observed in the pulmonary artery.² One conclusion that can be drawn from this observation is that acute clinical symptoms are

critical dependent upon the location of the bubbles. Bubbles in the brain, for instance, could give few symptoms, as large areas of the brain are clinically silent. Bubbles in joints, on the other hand, would give symptoms, because of the rich innervation by pain receptors in these areas. One effect of this would be that we have to distinguish between primary and secondary effects of bubbles. The primary effects are related to the mechanical effect of the bubbles, which may be blockage of the circulation or distortion of tissue. The secondary effects are related to the numerous effects of the bubble surface, with activation of a large number of biochemical and cellular mechanisms. It seems obvious that this secondary effect can occur without any acute signs or symptoms.

When do bubbles form ?

Most, if not all, practical decompressions will lead to some degree of gas bubble formation in the organism

and the risk of decompression illness (DCI). The predominant theory about the growth of bubbles is that bubbles grow from preformed nuclei, as the resistance of “pure solutions” to supersaturation and gas phase development is considerable.³ One likely theory is that the nuclei are composed of small (approximately 1 micron) stable gas bubbles.⁴

The exact threshold for bubble formation is not known, but it is probably in the range of 50-70 kPa in tissue⁵ and even lower in the vascular system. Eckenhoff et al. demonstrated that saturation at 3.7 msw on air was sufficient to produce bubbles in the pulmonary artery in man.⁶ The conclusion from this study must be that gas bubbles will form in the vascular system at any supersaturation and that the concept of a minimum tolerable limit of supersaturation, as least for the venous system, may only relate to clinical symptoms and not to bubble formation. Adding to this problem is the fact that it has been demonstrated repeatedly that large inter- and intra-individual differences in bubble forming “ability” exist. Factors like sex, age, body build, circulation, temperature, blood composition and degree of exercise seem to play a role.^{7,8} Cavitation in joints, for example, has been demonstrated without any supersaturation following violent movements. Even under experimental conditions where the circulation is kept stable, a variation in vascular bubbles between individuals, often a factor of 10, following decompression is observed.⁹ Furthermore, there are data indicating that there is a large difference in susceptibility to decompression sickness that is not directly related to the amount of vascular gas bubbles observed.¹⁰

Where do bubbles form ?

Bubbles have been observed in many tissues in the body following decompression. They are most commonly found in the vascular system, the white matter of the CNS, in abdominal fat, in synovial fluids and in muscles.¹¹ Following quite severe decompressions, we did not observe bubbles in the muscles themselves, but only on fascia.

Harvey et al. studied the limits for vascular bubble formation in cats, both at rest and after electrical stimulation and tissue injury.¹² The conclusion from these studies was that at marginal exposures, stimulation or injury was needed for bubble formation. At higher supersaturations, bubbles occurred at rest, the time of occurrence determined by the fat content. Essentially the same results have been obtained in frogs and rats.¹³ Based on these studies, the authors concluded that gas bubbles are chiefly intravascular and that they are responsible for nearly all important phases of the syndrome of decompression sickness. Only in very severe cases did extravascular bubbles play a role and then only in lipid rich structures.

Venous bubbles

There is evidence from many studies that gas bubbles occur in the venous system during most decompressions.^{14,15} Several studies have documented the relationship between the occurrence of many venous bubbles and the risk for clinical symptoms requiring treatment.^{16,17} This, together with the fact that bubbles probably are present in the venous system during most decompressions, suggests that a diver complaining of pain in a joint may be suffering from two different conditions, namely tissue gas in and around the joint and pulmonary gas embolism.

Arterial bubbles

Gas bubbles in the arteries have been detected in divers after excursions,¹⁸ during decompression from saturation dives¹⁹ and at autopsy after fatal accidents.²⁰ Arterial gas bubbles have also been observed in large animals during and after decompression.²¹⁻²³ Thus, there is no doubt that arterial gas bubbles occur during or after some decompressions.

In divers, there are several possible pathways by which venous bubbles may reach the arterial circulation. First, venous gas bubbles may travel through the pulmonary circulation and enter the pulmonary veins and the left atrium, although the pulmonary circulation is usually considered to be a good filter for gas bubbles as well as for other emboli. Second, venous gas bubbles may pass through a patent foramen ovale (PFO) or other extraordinary connections in the heart to reach the left side of the heart. Third, if the lung has been overinflated during a rapid ascent, gas may escape directly into the pulmonary veins after alveolar rupture.²⁴

Finally, gas bubbles may form in the arterial circulation if the decompression rate is sufficiently fast >0.3-1 fsw/sec.²⁵ All gas nuclei in the blood will not be destroyed at compression and supersaturation of the arterial blood may occur during the rapid decompression. However, an experimental study using goats did not succeed in demonstrating such bubbles in the arterial circulation after a short hyperbaric exposure and a rapid decompression.²⁶

In as many as 20-34% of humans, dependent on age, the foramen ovale is patent after foetal life.²⁷ Normally it is functionally closed, since the pressure in the left atrium is higher than the pressure in the right atrium and the septum primum functions as a valve. However, a spontaneous shunt, not dependent on a Valsalva manoeuvre or other factors to change the pressure gradient between the atria, is diagnosed in 5-6% of humans using contrast echocardiography.^{28,29}

Tissue bubbles

It seems reasonable to assume that tissue bubbles can occur if the gas load is high enough. This was apparently shown for the spinal cord by Francis et al., who showed tissue bubbles in the white matter following rapid decompression after a 15 minute dive to 300 fsw on air.³⁰ This work has, however, recently been challenged by Palmer, who claims that all changes observed could be explained by gas bubbles inside vessels.³¹ This is also supported by the recent work of Sharpe and Broome, who showed that there was no relationship between the fat content of the spinal cord and the occurrence of gas bubbles.³² Even if the exact mechanism of tissue injury is controversial, the evidence seems to indicate that vascular processes are the more important ones.

Primary bubble effects

Initially, the bubbles will lead to changes, mainly due to their direct mechanical effects. When gas bubbles form and expand they can obstruct the arterial and venous circulation, leading to tissue ischaemia, or they can damage the tissue and induce pain by direct pressure effects. Bubble formation may influence circulation by mechanical obstruction. Venous obstruction may lead to oedema and arterial obstruction may lead to tissue ischaemia, both of which have been observed after decompression.³³ This can reduce gas elimination, both by increasing diffusion distances and by reducing blood flow.

One important primary effect of the bubbles, which is often forgotten, is the reduction in gas elimination caused by these bubbles. Both theoretical³⁴ and experimental³⁵ studies have demonstrated that gas bubbles in the tissue will increase gas elimination time. In another study, it was shown that bubbles significantly increased the time constant of the slow component of the bi-exponential curve describing the nitrogen concentration in the pulmonary artery.³⁶ This can be seen in Figure 1.

Obstruction of flow to the tissue by bubbles may increase elimination time even more. Computer simulations have shown that a high number of bubbles can increase the time constant of gas elimination from muscles from 50 to 2,000 minutes (Flook, Personal communication 1999). This is partly taken into account by the new US Navy (USN) diving tables, where gas elimination is considered to be linear, not exponential.³⁷ However, in reality the problem may be even more complex as bubbles in the circulation may increase the transport of gas to the lungs.³⁸

Secondary bubble effects

When gas bubbles are formed protein denaturation takes place at the blood-gas interface.³⁹ The gas-blood




Figure 1. The relationship between the number of bubbles in the pulmonary artery and the time constant for the elimination of nitrogen from the pulmonary artery. Mean values with 95% confidence intervals.

interface is a thin layer, approximately 20 nm thick, consisting of fibrin and gamma globulin.^{40,41} This layer acts as a foreign substance, activating formed elements of the blood and inducing biochemical changes such as complement activation.⁴² These mechanisms play a significant role in the response of the tissue to venous gas emboli and are probably the basis for any long term effects that may occur. At present, our understanding of the importance of these mechanisms is quite fragmentary. Much further work is needed to determine the relative influence of mechanical and biochemical effects of gas embolism. An understanding of these processes is necessary for development of a rational approach to treating or preventing injury caused by vascular bubbles.

Inflammation

The response of an organism to injury is termed inflammation. In decompression sickness this inflammatory process is initiated by the surface of the bubbles, which is regarded by the organism as a foreign substance. However, this process can also be initiated by direct mechanical injury to the tissue. One important mediator of the inflammatory process is activation of the complement system. Gas bubbles activate the complement system *in-vitro*.⁴³ The degree of activation is dependent upon the amount of gas infused, varies considerably over several months in one individual and is not dependent upon the gas composition of the bubble. No relationship was seen between the degree of C5a activation *in vitro* and the level of C5a observed *in vivo* after air dives.⁴⁴

Leucocytes are involved in many aspects of tissue injury and inflammation. Several studies have documented that leucocytes are activated by decompression. Philp et al.⁴⁵ showed that decompression led to a reduction in both

thrombocytes and leucocytes. These reductions seem to be related to the presence of gas bubbles. During decompression from a saturation dive to 440 msw, where no gas bubbles could be detected, there was no reduction in leucocyte number.⁴⁶ In another saturation dive, Benestad et al.⁴⁷ showed that decompression lead to activation of neutrophils. Contrary to what has been observed in thrombocytes, no activation of leucocytes seems to occur in vitro at pressure.

Coagulation

Aggregation of thrombocytes may lead to the formation of blood clots, thus forming solid emboli which may compound the effect of gas bubbles in the lung. Thorsen et al.⁴⁸ showed that gas bubbles lead to aggregation of thrombocytes in-vitro. Aggregation in-vitro was strongest when the bubble diameter was between 40 and 120 μm . The degree of aggregation does not seem to be dependent upon the gas content of the bubble, but only on its surface properties.⁴⁹ Aggregation of thrombocytes by gas bubbles can be considerably enhanced by adrenaline (epinephrine).⁵⁰

Vasoconstriction/dilatation

Gas bubbles can induce vasoconstriction by direct effects on the vascular wall⁵¹ or vasodilation or constriction by initiating the release of different vasoactive substances.^{52,53}

Bubble effects on different organ systems

In the following the bubble effects will be described on some organ systems. Primary and secondary effects will be described together, as they often are quite difficult to distinguish.

Endothelial damage

Chryssanthou et al. have shown that animals exposed to decompression will show breakdown of the blood-brain barrier and the blood-lung barrier.⁵⁴ Broman et al. have demonstrated that even very short contact between gas bubbles and endothelium (1-2 minutes) will lead to such breakdown.⁵⁵ Furthermore, studies in rabbits indicate that such contact leads to endothelial damage and progressive reduction on cerebral blood flow and function. In a study by Smith et al.⁵⁶ endothelial damage could be demonstrated in pigs exposed to severe decompressions. We were able to demonstrate changes in the endothelium in pigs following exposure to gas bubbles. We found that these changes occurred at an exposure of approximately 1.5 bubbles/cm², equivalent to approximately Grade II-III on the Spencer scale.⁵⁷

Even minimal endothelial injury can induce activation of both biochemical and cellular responses, which could form the basis for tissue injury following decompression.

Lung function changes

Gas emboli may block some parts of the pulmonary vascular bed, reducing or preventing blood flow through the regions of lung served by those vessels. This leads to an initial rise in the pulmonary artery pressure (PAP) and pulmonary vascular resistance (PVR), and a decrease in arterial oxygen tension (P_{aO_2}).^{9,58,59} Following decompression, changes in diffusion capacity⁶⁰ and lung function changes similar to "small airways disease" have been seen.⁶¹

Central Nervous System changes.

Central nervous changes in DCI are probably caused by several mechanisms. In severe DCI, both vascular bubbles and in-vivo bubble formation probably plays a role.³³

Exposure to vascular bubbles without clinical symptoms do not seem to have a serious effect upon the spinal cord.⁶² In this group of 10 amateur and 10 professional divers, five of whom had suffered from DCI, no changes could be seen. In the brain, changes in the endothelial layer of the ventricles could be detected in a group of divers.⁶³ A possible explanation is that this damage is caused by gas bubbles in the spinal fluid, such bubbles will probably primarily adhere to the lining of the ventricles.

Numerous studies have shown that circulating gas bubbles change the blood-brain barrier, this is described above.

Bone

Aseptic bone necrosis is regarded as an occupational hazard for all workers under pressure.⁶⁴ There is clear indications that the incidence of bone necrosis is linked to decompression. This is perhaps best demonstrated by the fact that this disease, which is quite rare in the industrial world, is considerably more prevalent in the developing world, where diving practices produce a high incidence of decompression sickness.

Conclusions.

A recent consensus conference on long term health effects of diving¹¹ concluded that "changes can be seen in

lungs, CNS and bone in divers who had not had any decompression accidents", or perhaps more accurately, in divers who had not been treated for decompression illness. The mechanisms for these changes are not clear, but as bubbles in the vascular system have been observed frequently in divers, a reasonable working hypothesis would be that vascular bubbles may play a major role in this. There are still many details missing how the bubbles affect the organism. An understanding of the mechanisms involved is important, however, both for preventing injury and for treating the damage caused by these bubbles.

References

- 1 Behnke AR. Decompression sickness following exposure to high pressures. In *Decompression sickness*. Fulton JF. ed. Philadelphia: Saunders, 1951; 53-89
- 2 Spencer MP. Decompression limits for compressed air determined by ultrasonically detected blood bubbles. *J Appl Physiol* 1996; 40 (2): 229-235
- 3 Hemmingsen EA. Nucleation of bubbles in vitro and in vivo. In *Diving in animals and man*. Brubakk AO, Kanwisher J and Sundnes F. eds. Trondheim: Tapir Publishers, 1986, 43-59
- 4 Yount DE. Growth of bubbles from nuclei. In *Diving in animals and man*. Brubakk AO, Kanwisher J and Sundnes F. eds. Trondheim: Tapir Publishers, 1986, 131-164.
- 5 Daniels S. Ultrasonic monitoring of decompression procedures. *Phil Trans R Soc Lond* 1984; B304: 153-175
- 6 Eckenhoff RG, Olstad CS and Carrod G. Human dose-response relationship for decompression and endogenous bubble formation. *J Appl Physiol* 1990; 69: 914-918
- 7 Jones HB. Gas exchange and blood-tissue perfusion factors in various body tissues. In *Decompression sickness*. Fulton JF. ed. Philadelphia: Saunders, 1951; 279-321.
- 8 Vann RD. Exercise and circulation in the formation and growth of bubbles. In *Supersaturation and bubble formation in fluid and organisms*. Brubakk AO, Hemmingsen BB and Sundnes G. eds. Trondheim: Tapir Publishers, 1989; 235-258.
- 9 Vik A, Jenssen BM and Brubakk AO. Comparison of haemodynamic effects during venous air infusion and after decompression in pigs. *Eur J Appl Physiol* 1994; 68: 127-133
- 10 Ward CA, Weathersby PK, McCullough D and Fraser WD. Identification of individuals susceptible to decompression sickness. In *Underwater Physiology IX*. Bethesda, Maryland: UHMS, 1987; 239-247
- 11 Hope A, Lund T, Elliot DH, Halsey MJ and Wiig H. *Long term health effects of diving*. Bergen: NUTEC, 1994
- 12 Harvey EN. Animal experiments on bubble formation. Part I. Bubble formation in cats. In *Decompression sickness*. Fulton JF. ed. Philadelphia: Saunders, 1951; 115-144
- 13 Blinks LR, Twitty VC and Whitaker DM. Animal experiments on bubble formation. Part II. Bubble formation in frogs and rats. In *Decompression sickness*. Fulton JF. ed. Philadelphia: Saunders, 1951; 145-164.
- 14 Gardette B. Correlation between decompression sickness and circulating bubbles in 232 divers. *Undersea Biomed Res* 1979; 6: 99-107
- 15 Nishi RY. Doppler evaluation of decompression tables. In *Man in the sea. Vol I*. Lin YC and Shida KK. eds. San Pedro: Best Publishing Company, 1990; 297-316.
- 16 Nashimoto I and Gotoh Y. Relationship between precordial Doppler ultrasound records and decompression sickness. In *Underwater Physiology VI*. Schilling CW and Beckett MW. eds. Bethesda, Maryland: Federation of American Societies for Experimental Biology, 1978; 497-501
- 17 Nishi RY. Doppler and ultrasonic bubble detection. In *The physiology and medicine of diving. 4th Edition*. Bennett PB and Elliott DH. eds. London: WB Saunders Company, 1993; 433-453
- 18 Brubakk AO, Peterson R, Grip A, Holand B, Onarheim J, Segadal K, Kunkle T and Tønjum S. Gas bubbles in the circulation of divers after ascending excursions from 300 to 250 msw. *J Appl Physiol* 1986; 60: 45-51
- 19 Hjelle JO, Eatock BC, Nordahl SH and Dick AP. Doppler monitoring during 3 dives to 360 msw. In *XIIIrd Annual Meeting of the European Undersea Biomedical Society*. Marroni A and G. Oriani G. eds. Palermo: European Undersea Biomedical Society, 1987; 357-364
- 20 Haymaker W and Johnston AD. Pathology of decompression sickness: a comparison of lesions in airmen with those in caisson workers and divers. *Mil Med* 1955;117: 285-306
- 21 Bove AA, Hallenbeck JM and Elliot DH. Circulatory responses to venous air embolism and decompression sickness in dogs. *Undersea Biomed Res* 1974; 1: 297-220
- 22 Vik A, Brubakk AO, Hennessy TR, Jenssen BM, Ekker M and Slørdal S. Venous air embolism in swine: transport of gas bubbles through the pulmonary circulation. *J Appl Physiol* 1990; 69: 237-244
- 23 Vik A, Jenssen BM and Brubakk AO. Effect of aminophylline on transpulmonary passage of venous air emboli in pigs. *J Appl Physiol* 1991; 71: 1780-1786
- 24 Neuman TS and Bove AA. Combined arterial gas embolism and decompression sickness following no-stop dives. *Undersea Biomed Res* 1990; 17: 429-436
- 25 Kunkle TD and Beckman EC. Atraumatic air

- embolism in diving. *Undersea Biomed Res* 1981; 8: 11-12
- 26 Powell MR, Spencer MP, Smith MT, Beckmann EL and Kunkle TD. In situ arterial bubble formation and "atraumatic air embolism". *Undersea Biomed Res* 1982; 9 (Suppl): 10
- 27 Hagen PT, Scholz DG and Edwards WD. Incidence and size of patent foramen ovale during the first 10 decades of life: an autopsy study of 965 normal hearts. *Mayo Clin Proc* 1984; 59: 17-20
- 28 Guggiari M, Lechat PH, Garen-Colonne C, Fuscuardi J and Viars P. Early detection of patent foramen ovale by two-dimensional contrast echocardiography for prevention of paradoxical air embolism during sitting position. *Anesth Analg* 1988; 67: 192-194
- 29 Lynch JJ, Schuchard GH, Gross CM and Wann LS. Prevalence of right-to-left atrial shunting in a healthy population: detection by Valsalva maneuver contrast echocardiography. *Am J Cardiol* 1984; 53: 1478-1480
- 30 Francis TJR, Griffin JL, Homer LD, Pezeshkpour GH, Dutka AJ and Flynn ET. Bubble-induced dysfunction in acute spinal cord decompression sickness. *J Appl Physiol* 1990; 68: 1368-1375
- 31 Palmer AC. Nature and incidence of bubbles in the spinal cord of decompressed goats. *Undersea Hyper Med* 1997; 24: 193-200
- 32 Broome JR and Sharpe RP. Spinal cord lipid levels in a porcine model of spinal cord decompression sickness. *Undersea Hyper Med* 1997; 24: 323-328
- 33 Francis TJR and Gorman DF. Pathogenesis of the decompression disorders. In *The physiology and medicine of diving. 4th Edition.* Bennett PB and Elliott DH. eds. London: WB Saunders Company, 1993; 454-480
- 34 Van Liew HD and Burkard ME. Density of decompression bubbles and competition for gas among bubbles, tissue, and blood. *J Appl Physiol* 1993; 75: 2293-2301
- 35 D'Aoust BG, Smith KH and Swanson HT. Decompression-induced decrease in nitrogen elimination rate in awake dogs. *J Appl Physiol* 1976; 41: 348-355
- 36 Flook V, Brubakk AO, Eftedal O, Holmen I, Ustad A-L and Koteng S. *The effect of oxygen and of decompression bubbles on inert gas washout. STF23 A94031* Trondheim: Sintef, 1994
- 37 Thalmann ED, Parker EC, Survanshi S and Weathersby PK. Improved probabilistic decompression model risk predictions using linear-exponential kinetics. *Undersea Hyper Med* 1997; 24: 255-274
- 38 Kindwall EP, Baz A, Lightfoot EN, Lanphier EH and Seireg A. Nitrogen elimination in man during decompression. *Undersea Biomed Res* 1975; 2: 285-297
- 39 Lee WH and Hairston P. Structural effects on blood proteins at the gas-blood interface. *Fed Proc* 1971; 30: 1615-1622
- 40 Lehto V-P, Kantola J, Tervo T and Laitinen LA. Ruthenium red staining of blood-bubble interface in acute decompression sickness in rat. *Undersea Biomed Res* 1981; 8: 101-111
- 41 Vroman L, Adams AL and Klings M. Interactions among human blood proteins at interfaces. *Fed Proc* 1971; 30: 1494-1502
- 42 Tamiya T, Yamasaki M, Maeo Y, Yamashiro T, Ogoshi S and Fujimoto S. Complement activation in cardiopulmonary bypass, with special reference to anaphylatoxin production in membrane and bubble oxygenators. *Ann Thorac Surg* 1988; 46: 47-57
- 43 Bergh K, Hjelde A, Iversen OA and Brubakk AO. Variability over time of complement activation induced by air bubbles in human and rabbit sera. *J Appl Physiol* 1993; 74: 1811-1815
- 44 Hjelde A, Bergh K, Brubakk AO and Iversen OJ. Complement activation in divers after repeated air/heliox dives and its possible relevance to DCS. *J Appl Physiol* 1995; 78: 1140-1144, 1995.
- 45 Philp R, Inwood M and Warren B. Interactions between gas bubbles and components of the blood: Implications in decompression sickness. *Aerospace Med* 1972; 43: 946-953
- 46 Shirakawa Y, Suzuki S, Hashimoto A and Oiwa H. Effects of deep saturation diving on the lymphocyte subsets of healthy divers. *Undersea Hyper Med* 1994; 21: 277-286
- 47 Benestad HB, Hersleth IB, Hardersen H and Molvær OI. Functional capacity of neutrophil granulocytes in deep-sea divers. *Scand J Clin Lab Invest* 1990; 50: 9-18
- 48 Thorsen T, Brubakk AO, Øvstedal T, Farstad M and Holmsen H. A method for production of N₂ microbubbles in platelet-rich plasma in an aggregometer-like apparatus, and effect of platelet density in vitro. *Undersea Biomed Res* 1986; 13: 271-288
- 49 Thorsen T, Klausen H, Lie RT and Holmsen H. Bubble-induced, aggregation of platelets: effects of gas species, proteins and decompression. *Undersea Hyper Med* 1993; 20: 101-120
- 50 Thorsen T, Ovstedal T, Vereide A and Holmsen H. Effect of platelet agonists on the reduction in platelet density caused by microbubbles in vitro. *Undersea Biomed Res* 1986; 13: 289-303
- 51 Malik AB. Pulmonary Microembolism. *Physiological Reviews* 1983; 63: 1114-1207
- 52 Garcia-Szabo RR, Minnear FL, Bizios R, Johnsen A and Malik AB. Role of thromboxane in the pulmonary response to pulmonary microembolism. *Chest* 1983; 83: 768-788
- 53 Malik AB and Johnson A. Role of humoral mediators in the pulmonary vascular response to pulmonary embolism. In *Pulmonary vascular physiology and pathophysiology.* Weir K and Reeves JT. eds. New York and Basel: Macel Dekker Inc., 1995, 445-468
- 54 Chryssanthou C, Springer M and Lipschitz S. Blood-

- brain and blood-lung barrier alterations by dysbaric exposure. *Undersea Biomed Res* 1977; 4: 111-116
- 55 Broman T, Branemark PI, Johansson B and Steinwell O. Intravital and post-mortem studies on air embolism damage of the blood-brain-barrier. *Acta Neur Scand* 1996; 42: 146-152
- 56 Smith KH, Stegall PJ, Harker LA, Slichter SJ, Richmond VL, Hall MH and Huang TJ. Investigation of hematologic and pathologic response to decompression. #N00014-71-C-0273. Seattle: Virginia Mason Research Center, 1978
- 57 Nossum V and Brubakk AO. Endothelial damage by bubbles in the pulmonary artery of the pig. *Undersea Hyper Med* 1999; : 1-8, 1999
- 58 Verstappen FTJ, Bernards JA, and Kreuzer F. Effects of Pulmonary Gas Embolism on Circulation and Respiration in the Dog. I. Effects on Circulation. *Pflügers Arch* 1977; 368: 89-96
- 59 Verstappen FTJ, Bernards JA and Kreuzer F. Effects of Pulmonary Gas Embolism on Circulation and Respiration in the Dog. IV. Origin of Arterial Hypoxemia during Pulmonary Gas Embolism. *Pflügers Arch* 1977; 370: 71-75, 1977.
- 60 Thorsen E, Segadal K and Kambestad BK. Mechanisms of reduced pulmonary function after a saturation dive. *J Eur Respir* 1994; 7: 4-10
- 61 Thorsen E, Segadal K, Kambestad BK and Gulsvik A. Divers' lung function: small airways disease? *Brit J Industr Med* 1990; 47: 519-523
- 62 Mørk SJ, Morild I, Brubakk AO, Eidsvik S and Nyland H. A histopathologic and immunocytochemical study of the spinal cord in amateur and professional divers. *Undersea Hyper Med* 1994; 21: 391-402
- 63 Mørk SJ and Morild E. A neuropathological study of the ependymoventricular surface in divers brains. *Undersea Hyper Med* 1994; 21: 43-51
- 64 Evans A, King JD, McCallum RI, Thickett VB, Throwbridge WP and Walder DN. Aseptic bone necrosis in commercial divers: a report from the decompression sickness central registry and radiological panel. *Lancet* 1981; i: 384-388

Dr Alf O Brubakk was one of the guest speakers at the 1999 SPUMS Annual Scientific Meeting. His address is Department of Physiology and Biomedical Engineering, Medical Faculty, The Norwegian University of Science and Technology, Trondheim, Norway. Telephone +47-7359-1005. Fax +47-7359-8904. E-mail Alf.O.Brubakk@medisin.ntnu.no.

37 DECOMPRESSION SICKNESS CASES TREATED IN THE DEPARTMENT OF UNDERWATER AND HYPERBARIC MEDICINE, ISTANBUL FACULTY OF MEDICINE

Akin S Toklu, Samil Aktas, Salih Aydin and Maide Çimsit

Abstract

The time interval between the onset of decompression sickness (DCS) and recompression therapy, and the first aid with medical treatments applied before reaching a recompression facility, will affect the outcome of the recompression therapy.

In this study 37 DCS cases were evaluated to find out the time interval to the onset of DCS, the type of the disease and symptoms, delay to recompression treatment, medical treatments applied during transport, recompression treatment protocols performed and outcomes.

All the patients were male. Professional divers (32 or 86.5%) outnumbered sports divers (5 or 13.5%). In 20 cases (54.1%) onset of symptoms was within the first 10 minutes after the dive. In three men (8.1%) symptoms came on underwater. Numbness, tingling and back pain were the most frequent symptoms reported by the patients with Type II DCS. Complete recovery was achieved in 32 (86.5%) of the cases by recompression therapy combined with medical treatment. Rehabilitation was needed in 12 (32.4 %) of the cases.

Omitted decompression was the most frequent cause of DCS in our cases. Additional hyperbaric oxygen therapy needed in delayed cases is evidence of the importance of immediate transport and adjunctive medical treatments.

Key Words

Decompression illness, first aid, transport, treatment.

Introduction

The hyperbaric facilities in Turkey are mostly situated in Istanbul. The only Hyperbaric and Underwater Medicine Department in civilian universities is in the Istanbul Faculty of Medicine. The Fisheries Research Institute also has a hyperbaric chamber for treating divers in Bodrum, in Aegean Sea region. The Turkish Navy has three hyperbaric units. Besides these public facilities, all with multiplace chambers, hyperbaric oxygen therapy is performed in three private hyperbaric centres, in Istanbul. The three private hyperbaric centres, which all have multiplace chambers and one also has a monoplace, are free standing and mostly use hyperbaric oxygen therapy for indications other than diving related disease. In March