

PLA₂ INHIBITION BY LIGNOCAINE: IS IT CLINICALLY RELEVANT ?

Georg Petroianu and Ursula Helfrich

Key words

Decompression illness, drugs, treatment.

Abstract

The place of lignocaine administration for DCI treatment seems to be well established. The rationale for its use is a putative anti-inflammatory effect of the drug, most probably due to its ability to inhibit phospholipase A₂ (PLA₂). The purpose of the study was to quantify "in vitro" lignocaine's ability to inhibit this key enzyme and to elucidate the type of inhibition. Lignocaine inhibits PLA₂ through interaction with the enzyme-substrate complex. This occurs at plasma concentrations which are easily achievable clinically. Therefore the use of lignocaine as an anti-inflammatory drug seems warranted.

Introduction

The SPUMS Journal has published two papers on the use of lignocaine as adjuvant therapy in the treatment of decompression illness (DCI).^{1,2} While both authors agree that lignocaine has a well established place in DCI therapy and that the anti-inflammatory effect of lignocaine might be the strongest rationale for using it for this purpose, there appears to be little data available on the magnitude of these anti-inflammatory effects.

Lignocaine is a known phospholipase A₂ inhibitor.^{3,4} This study was to quantify "in vitro" lignocaine's ability to inhibit this key enzyme and to elucidate the type of inhibition.

Material and Methods

Blood samples were taken from nine healthy human volunteers. PLA₂ derived from the platelet membranes was incubated for 30 minutes with either TRIS buffer (native samples or controls) or lignocaine. Lignocaine concentrations of 1, 10 or 100 µg/ml (4.3; 43.0; 430 µM) were used. PLA₂ activity was measured by a modification of the method described by Flesch⁵ and Sundaram,⁶ while protein concentrations were determined by a modified Lowry method.^{7,8} PLA₂ activities were expressed in pmol/mg protein/min. Mean values were used for statistical analysis with the Mann-Whitney rank order test. Baseline values (native activity) were considered to be 100%. All other values were expressed as a percentage of the baseline value.

For K_M and V_{MAX} determinations commercially available purified porcine PLA₂ (Sigma; Steinheim, Germany) was incubated with different substrate concentrations (0-300 µM) in the presence or absence of lignocaine (100 µg/ml = 430 µM) for 30 minutes. The PLA₂ activity was determined in a commercially available radioactive PLA₂ assay (Scintillation Proximity Assay: SPA; Amersham, Braunschweig, Germany). Data were plotted as Michaelis-Menten and Lineweaver-Burk diagrams.

Results

Lignocaine inhibits human platelet membrane PLA₂ activity in a statistically significant manner. However in the concentration range used (1-100 µg/ml) no dose dependency could be observed: the lowest concentration used led to a maximal inhibition of the enzyme (Figure 1).




Figure 1. Lignocaine inhibits human platelet membrane PLA₂ activity in a statistically significant manner ($p \geq 0.010$). However in the concentration range used (1 - 100 µg/ml) no dose dependency could be observed.

Lineweaver-Burk representation of the data (using porcine PLA₂) suggests an interaction of lignocaine with the PLA₂ molecule and the enzyme-substrate-complex (non-competitive or mixed inhibition). The coordinates of the intersection point are $x = -0.16$ and $y = -0.06$. The inhibitor constants K_I (for the enzyme-inhibitor; EI) and K_I' (for the enzyme-substrate-inhibitor; ESI) were calculated. K_I (4,800 µM) is one order of magnitude higher than K_I' (409 µM) suggesting that the main mode of action of lignocaine is interference with the enzyme-substrate complex formation. The correlation coefficient for data determined in the absence of the inhibitor is $r_{\text{native}} = 0.96$ and for data determined in the presence of the inhibitor is $r_{\text{Lignocaine}} = 0.98$ (See Figure 2 on page 10).




Figure 2. Lineweaver-Burk representation of the data (porcine PLA₂) suggests an interaction of lignocaine with the PLA₂ molecule and the enzyme-substrate-complex [non-competitive (mixed) inhibition].

Discussion

The effective plasma concentration range of lignocaine in humans is 1-20 μg/ml (4-80 μM). The lowest lignocaine concentration used (1 μg/ml) produced maximal inhibition of the human platelet derived PLA₂. Therefore the anti-inflammatory effect of lignocaine is easily achievable using common clinical dosages. The data derived from experiments using porcine enzyme show that the anti-inflammatory effect of lignocaine is mainly due to interaction with the enzyme-substrate-complex. The inhibitory constant K_I for porcine PLA₂ is in the 400 μM range. The most probable explanation for this value (five times higher than the upper limit of the effective plasma concentration range) is the higher sensitivity of the human enzyme to lignocaine inhibition compared with the porcine variant. Different activities/sensitivities for PLA₂ of different origins are well recognised.⁹

Conclusion

We conclude that lignocaine's ability to inhibit PLA₂ through interaction with the enzyme-substrate-complex occurs at plasma concentrations which are easily achievable clinically. As such the use of lignocaine as an anti-inflammatory drug seems warranted.

References

- 1 Mitchell S. The role of lignocaine in the treatment of decompression illness: a review of the literature. *SPUMS J* 1995; 25 (4): 182-194
- 2 Cosh D. Lignocaine and non-steroidal anti-inflammatory drugs as adjuvant therapy in treating

decompression illness. *SPUMS J* 1998; 28 (2): 97-99

- 3 Denson DD, Worrell RT and Eaton DC. A possible role for phospholipase A₂ in the action of general anesthetics. *Am J Physiol* 1996; 270: C636-C644
- 4 Kunze H, Nahas N and Traynor JR. Effects of local anaesthetics on phospholipases. *Biochem Biophys Acta* 1976; 441: 93-102
- 5 Flesch I, Schmidt B and Ferber E. Acylchain specificity, and kinetic properties of phospholipase A₁, and A₂ of bone marrow derived macrophages. *Naturforsch* 1985; 40: 356-363
- 6 Sundaram GS, Shakir KMM, Barnes G and Margolis S. Release of phospholipase A₂, and triglyceride lipase from rat liver. *Biol Chem* 1978; 253: 7703-7710
- 7 Lowry OH, Rosebrough NJ, Farr AL and Randall RJ. Protein measurement with the folin phenol reagent. *Biol Chem* 1951; 193: 265-275
- 8 Petroianu G, Helfrich U, Schmitt A, Bergler W and Ruefer R. Dose-dependent inhibition of phospholipase A₂ by paraoxon *in vitro*: preliminary results. *J Appl Toxicol* 1997; 17: 421-424
- 9 Brockerhoff H and Jensen RG. *Lipolytic enzymes*. New York: Academic Press, 1974; 206-215

Dr Georg Petroianu is a Privat Dozent in the Department of Pharmacology and Toxicology, University of Heidelberg at Mannheim, Maybach Street 14-16, 68169 Mannheim, Germany. Phone +49-621-330-0328. Fax +49-621-339-0333. E-mail petroia@rumms.uni-mannheim.de

Ursula Helfrich is a medical student at the same university.

THE
SPUMS

HOME PAGE,

WHICH GIVES ACCESS TO THE

**SPUMS JOURNAL INDEX 1971-1998
 IS AT**

<http://www.SPUMS.org.au>